

INFALL REGIONS AND SCALING RELATIONS OF X-RAY SELECTED GROUPS

KENNETH RINES^{1,2} AND ANTONALDO DIAFERIO^{3,4}

Draft version February 3, 2010

ABSTRACT

We use the Fifth Data Release of the Sloan Digital Sky Survey to study X-ray-selected galaxy groups and compare their properties to clusters. We search for infall patterns around the groups and use these to measure group mass profiles to large radii. In previous work, we analyzed infall patterns for an X-ray-selected sample of 72 clusters from the ROSAT All-Sky Survey. Here, we extend this approach to a sample of systems with smaller X-ray fluxes selected from the 400 Square Degree serendipitous survey of clusters and groups in ROSAT pointed observations. We identify 16 groups with SDSS DR5 spectroscopy, search for infall patterns, and compute mass profiles out to 2-6 h^{-1} Mpc from the group centers with the caustic technique. No other mass estimation methods are currently available at such large radii for these low-mass groups, because the virial estimate requires dynamical equilibrium and the gravitational lensing signal is too weak. Despite the small masses of these groups, most display recognizable infall patterns. We use caustic and virial mass estimates to measure the scaling relations between different observables, extending these relations to smaller fluxes and luminosities than many previous surveys. Close inspection reveals that three of the groups are subclusters in the outskirts of larger clusters. A fourth group is apparently undergoing a group-group merger. These four merging groups represent the most extreme outliers in the scaling relations. Excluding these groups, we find $L_X \propto \sigma_p^{3.4 \pm 1.6}$, consistent with previous determinations for both clusters and groups. Understanding cluster and group scaling relations is crucial for measuring cosmological parameters from clusters. The complex environments of our group sample reinforce the idea that great care must be taken in determining the properties of low-mass clusters and groups.

Subject headings: galaxies: clusters: individual — galaxies: kinematics and dynamics — cosmology: observations

1. INTRODUCTION

A large fraction of all galaxies are members of groups. Groups are a less extreme and much more common type of system than galaxy clusters. Because they are less massive than clusters, groups are currently less well understood than clusters. Here, we investigate the optical properties of a sample of X-ray-selected groups to determine whether they can be modeled as scaled-down versions of clusters⁵. In particular, we study the outskirts of the groups to determine if their infall regions are readily identifiable as they are in the outskirts of clusters. With large samples of spectroscopic members, we then estimate the virial masses of the groups and determine if the groups obey the same scaling relations as clusters.

Galaxy clusters and groups are surrounded by infall regions in which galaxies are bound to the system but are not in equilibrium. The Cluster Infall Regions in the Sloan Digital Sky Survey (Rines & Diaferio 2006, hereafter CIRS) project showed that X-ray-selected clusters display a characteristic trumpet-shaped pattern in radius-redshift phase space diagrams. These patterns,

termed caustics, were first predicted for simple spherical infall onto clusters (Kaiser 1987; Regös & Geller 1989), but later work showed that these patterns reflect the dynamics of the infall region (Diaferio & Geller 1997, hereafter DG) and (Diaferio 1999, hereafter D99). CIRS and earlier similar studies (e.g., Geller et al. 1999; Rines et al. 2003) showed that the amplitude of the caustics yields an estimate of cluster mass profiles consistent with both virial and X-ray mass estimates where the techniques overlap. More recently, Diaferio et al. (2005) and Lemze et al. (2009) showed that caustic masses also agree well with mass estimates based on gravitational lensing. Because neither galaxies nor gas is expected to be in equilibrium outside the virial radius of a cluster, the caustic technique and weak lensing are the only well-studied methods for determining cluster mass profiles at large radii (see Diaferio 2009, for a recent review of the caustic technique).

CIRS showed that infall patterns are ubiquitous in nearby massive clusters selected by X-ray emission. The CIRS clusters are fairly massive clusters and generally have little surrounding large-scale structure (but see Rines et al. 2001, 2002). One might suspect that the presence of infall patterns is limited to massive, isolated clusters. However, other investigators have found infall patterns around the Fornax Cluster (Drinkwater et al. 2001), the Shapley Supercluster (Reisenegger et al. 2000), an ensemble cluster comprised of poor clusters in the Two Degree Field Galaxy Redshift Survey (Biviano & Girardi 2003), and even the galaxy group associated with NGC 5846 (Mahdavi et al. 2005). Here, we extend the study of infall patterns by studying a large number of systems

Electronic address: krines@cfa.harvard.edu

¹ Smithsonian Astrophysical Observatory, 60 Garden St, MS 20, Cambridge, MA 02138; krines@cfa.harvard.edu

² Department of Physics and Astronomy, Western Washington University, Bellingham, WA

³ Università degli Studi di Torino, Dipartimento di Fisica Generale “Amedeo Avogadro”, Torino, Italy; diaferio@ph.unito.it

⁴ Istituto Nazionale di Fisica Nucleare (INFN), Sezione di Torino, Torino, Italy

⁵ The division between groups and clusters is somewhat arbitrary; here, we define groups to be systems of galaxies with virial masses smaller than $10^{14} h^{-1} M_\odot$.

with smaller X-ray fluxes than the CIRS clusters. Because these systems typically have smaller masses and fewer member galaxies (more typical of groups), infall patterns might not be identifiable in these systems.

In particular, we use the new 400 Square Degree (400d) survey of clusters and groups in *ROSAT* PSPC pointed observations (Burenin et al. 2007). The 400d survey is the largest area cluster survey extending to flux limits of $\approx 1.4 \times 10^{-13} \text{ erg s}^{-1} \text{ cm}^{-2}$. This flux limit is a factor of 20 smaller than the flux limit of catalogs based on the *ROSAT* All-Sky Survey (RASS; Voges et al. 1999). We search for 400d clusters and groups in the spectroscopic footprint of the Sloan Digital Sky Survey Data Release 5 (Adelman-McCarthy et al. 2007, SDSS DR5). This approach is similar to CIRS and to the RASS-SDSS (Popesso et al. 2004, 2005) analysis of clusters in DR2. Compared to both the CIRS and RASS-SDSS samples, the 400d groups are expected to have significantly smaller masses on average.

One motivation for this study is to improve our understanding of cluster scaling relations, in particular to extend these relations to the regime of high-mass groups. Ambitious cluster surveys like the South Pole Telescope and Atacama Cosmology Telescope require a good understanding of cluster scaling relations to accurately measure cosmological parameters from the abundance and evolution of clusters. Quantifying the scatter in these relations and any Malmquist-like bias is crucial for obtaining robust cosmological constraints from cluster surveys (e.g., Stanek et al. 2006). Many forecasts for dark energy constraints from cluster surveys assume that scatter in the mass-observable scaling relations has a log-normal distribution (e.g., Mantz et al. 2008). Deviations from this assumption could significantly impact the constraining power of these surveys.

Recent studies of scaling relations that extend into the group regime have often reached contradictory conclusions. For instance, a common scaling relation is the $L_X - \sigma_p$ relation between the X-ray luminosity L_X of a cluster or group and its projected velocity dispersion σ_p . For massive clusters, this relation is typically found to have a steep slope of $4.4^{+0.7}_{-0.3}$ (Mahdavi & Geller 2001) or 3.7 ± 0.3 (Popesso et al. 2005). For groups, various studies have found slopes as shallow as 0.37 (Mahdavi et al. 2000, the fit is a broken power-law with a slope of 4.02 for clusters) and as steep as 4.7 ± 0.9 (Helsdon & Ponman 2000). A detailed optical-X-ray study by Osmond & Ponman (2004) recently found a slope of 2.5 ± 0.4 . The variety in slopes is produced by many factors, including differing definitions of L_X , and in some cases, possible evolution of the galaxies via dynamical friction (Helsdon et al. 2005). One problem is that few large, complete group catalogs are available, so many existing studies utilize either heterogeneous samples or include many groups with limited data on each group. One solution is to utilize X-ray selection (e.g., Mulchaey & Zabludoff 1998; Mahdavi et al. 2000), but existing cluster/group catalogs based on RASS (Popesso et al. 2005, CIRS) have been restricted to systems with relatively high X-ray fluxes (and therefore including few groups). The fainter X-ray flux limits of the 400d survey allow detection of several groups. Combining this catalog with optical data from the Sloan Digital Sky Survey (Stoughton et al. 2002), we

can test whether the scaling relations show similar behavior for systems with smaller fluxes more typical of galaxy groups. A complementary approach (that we do not apply here) to understanding scaling relations for low-mass systems is to identify the systems optically and use stacking analysis to measure their ensemble properties. This approach has been applied previously with RASS data (Dai et al. 2007; Rykoff et al. 2008).

We describe the data and the group sample in § 2. In § 3, we review the caustic technique and use it to estimate the group mass profiles, discuss cluster scaling relations, and compare the caustic mass profiles to simple parametric models. We discuss some individual groups in § 4. We compare the caustic mass profiles to X-ray and virial mass estimators in § 5. We discuss our results and conclude in § 6. We assume $H_0 = 100 h \text{ km s}^{-1} h^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$ throughout.

2. THE 400D-SDSS GROUP SAMPLE

2.1. Sloan Digital Sky Survey

The Sloan Digital Sky Survey (SDSS, Stoughton et al. 2002) is a wide-area photometric and spectroscopic survey at high Galactic latitudes. The Fifth Data Release (DR5) of SDSS includes 8000 square degrees of imaging data and 5740 square degrees of spectroscopic data (Adelman-McCarthy et al. 2007).

The spectroscopic limit of the main galaxy sample of SDSS is $r=17.77$ after correcting for Galactic extinction (Strauss et al. 2002). CAIRNS and CIRS found that infall patterns were detectable in clusters sampled to about $M^* + 1$, or $z \lesssim 0.1$ for SDSS data. For the current sample, the X-ray fluxes are much smaller than those of CAIRNS or CIRS, so these groups may not be as well sampled. If the 400d groups are less massive on average, they should contain fewer luminous galaxies than the CAIRNS and CIRS clusters. We therefore expect that infall patterns may be less common and/or poorly sampled in the 400d groups.

Note that SDSS is $\sim 85\text{--}90\%$ complete to the nominal spectroscopic limit. The survey has $\approx 7\%$ incompleteness due to fiber collisions (Strauss et al. 2002), which are more likely to occur in dense cluster fields. Because the target selected in a fiber collision is determined randomly, this incompleteness can theoretically be corrected for in later analysis. From a comparison of SDSS with the Millennium Galaxy Catalogue, Cross et al. (2004) conclude that there is an additional incompleteness of $\sim 7\%$ due to galaxies misclassified as stars or otherwise missed by the SDSS photometric pipeline. For our purposes, the incompleteness is not important provided sufficient numbers of group galaxies do have spectra.

2.2. X-ray Surveys for Clusters and Groups

In CIRS, we studied clusters contained in catalogs based on the *ROSAT* All-Sky Survey (RASS Voges et al. 1999). RASS is a shallow survey but it is sufficiently deep to include nearby, massive clusters and groups. RASS covers virtually the entire sky and is thus the most complete X-ray cluster survey for nearby clusters and groups. The flux limits of RASS-based surveys are $\approx 3 \times 10^{-12} \text{ erg s}^{-1} \text{ cm}^{-2}$ (Ebeling et al. 1998; Böhringer et al. 2000a).

The wide field-of-view of the *ROSAT* PSPC instru-

ment provides a large area for serendipitous discovery of sources in pointed observations. Several cluster catalogs have been created with this goal. The largest of these catalogs is the 400d survey, which contains essentially all pointed observations suitable for extragalactic surveys (Burenin et al. 2007), yielding a total survey of about 400 square degrees. Extended X-ray sources are detected with a wavelet technique and fluxes are measured in the ROSAT band (0.1-2.4 keV). The flux limit of the 400d survey is $\approx 1.4 \times 10^{-13} \text{ erg s}^{-1} \text{ cm}^{-2}$, a factor of ~ 20 fainter than the RASS-based catalogs. The effective area of the survey depends on the flux limit and is calibrated with simulated observations of real ROSAT cluster data. The detection algorithm is optimized for luminous clusters ($L_X > 5 \times 10^{43} \text{ erg s}^{-1}$ at moderate redshift ($z > 0.3$)). The groups studied here have smaller X-ray luminosities and are thus expected to be physically smaller than higher-luminosity clusters, but they are also at much lower redshifts, so their angular sizes are not too different from the luminous clusters at $z > 0.3$. The ROSAT observations are typically not deep enough to separate diffuse group emission from contributions from individual galaxy halos.

We restrict our analysis to clusters and groups with $z \leq 0.10$ (§2.1). Our sample contains 16 groups within the SDSS DR5 spectroscopic footprint. We will refer to this sample as the 400d-SDSS groups hereafter. Figure 1 shows the X-ray luminosities of these systems compared to CIRS. The range of L_X in the two surveys is similar, but the 400d-SDSS sample contains many more systems with small L_X than does CIRS.

3. RESULTS

3.1. Infall Patterns around X-ray Groups

We first search for well-defined infall patterns around the 400d-SDSS groups. Analogously to CIRS, we plotted radius-redshift diagrams for all groups in the 400d X-ray catalog covered by DR5 with $z < 0.10$. We assign a “by-eye” classification of each group’s infall pattern: “clean” for groups with few background and foreground galaxies, “intermediate” for groups with apparent infall patterns but significant contamination from either related large-scale structure or foreground and background objects, and “weak” for groups with little apparent infall pattern. CIRS successfully applied the caustic technique to clusters classified as “intermediate” or “weak”; the classification scheme is thus fairly conservative. We use this classification scheme *only* to show the dependence of the infall pattern appearance on group mass (using L_X as a proxy) and the sampling depth.

Figure 1 shows the dependence of this subjective classification on X-ray luminosity and redshift. The 400d-SDSS sample contains 16 groups; six contain “clean” infall patterns and three (19%) show weak infall patterns. The percentage in the last category is larger than for the CIRS clusters (4%), although the significance of this difference is difficult to assess (the 400d-SDSS sample is small and the categories are not robustly defined).

Figure 1 demonstrates the expanded parameter space covered by the 400d-SDSS groups compared to the CIRS sample. In particular, the 400d-SDSS sample includes many more systems with $L_X \sim 10^{42} \text{ h}^{-2} \text{ erg s}^{-1}$ than CIRS. Table 1 lists the groups in the 400d-SDSS




FIG. 1.— Redshift versus X-ray luminosity (0.1-2.4 keV) for X-ray clusters from CIRS (small squares) and 400d-SDSS groups (large hexagons and stars) contained in the SDSS DR5 spectroscopic survey region. Filled hexagons, open hexagons and six-pointed stars indicate 400d-SDSS groups with “clean”, “intermediate”, and “weak” infall patterns respectively. These classifications for CIRS clusters are represented by filled squares, open squares, and crosses. Large open circles around some of the 400d-SDSS groups indicate systems contaminated by nearby structure (see §3.3). The dashed line and lower solid lines show flux limits of $f_X > 1.4$ and $3 \times 10^{-13} \text{ erg s}^{-1}$ respectively; the effective survey areas at these limits for the 400d survey are 232 and 370 square degrees respectively (not all of this area is contained within the SDSS DR5 spectroscopic footprint). The upper solid line shows the flux and redshift limits of the CIRS cluster sample. The dash-dotted line shows the redshift and luminosity limits of CAIRNS.

sample, their X-ray positions and luminosities, their central redshifts and velocity dispersions (see below), and the projected radius R_{comp} within which the SDSS DR5 spectroscopic survey provides complete spatial coverage. For several groups, the caustic pattern disappears beyond R_{comp} because of this edge effect. Figures 2-3 show the infall patterns for the 400d-SDSS sample.

Another factor which determines the presence or absence of a “clean” infall pattern is the surrounding large-scale structure. For example, the redshift-radius diagrams of groups within the infall regions of more massive clusters reflect the kinematics of the cluster’s infall region (e.g., A2199, see Rines et al. 2002). Closer inspection of the 400d-SDSS groups reveals that the caustic technique misidentifies four of the 16 groups. That is, the galaxy group associated with the 400d X-ray source is embedded in a larger system containing more galaxies. The hierarchical clustering algorithm used by the caustic technique to define the center of the system then mistakes the center of the larger system as the center of the 400d group. Ignoring these four groups, the percentage of “weak” infall patterns increases from 19% to 25% (three of the 12 remaining 400d-SDSS groups). Although the statistics are poor, this result suggests that the 400d-SDSS groups might have less well-defined infall patterns than the generally more massive CIRS clusters.

We attempt to quantify this difference more robustly by defining a simple statistic to quantify the contrast of the infall pattern with respect to the local background. We define the contrast C_{200} to be the ratio of the number




FIG. 2.— Redshift versus radius for SDSS galaxies around the first eight X-ray groups in the 400d-SDSS sample. The caustic pattern is evident as the trumpet-shaped regions with high density. The solid lines indicate our estimate of the location of the caustics in each group. Vertical lines in each panel indicate the radius R_{comp} where the spatial coverage of the SDSS DR5 spectroscopic survey is no longer complete (no line is shown if $R_{comp} > 7h^{-1}\text{Mpc}$). Squares around group names indicate systems with contamination from nearby groups or clusters (see §3.3). Figure 3 shows similar plots for the rest of the sample.

of galaxies within projected radius r_{200} within the caustics N_{mem} to the number of “near-background” galaxies N_{bgd} , those within $\pm 5\sigma_{200}$ of the cluster redshift but outside the caustics. The radius r_{200} is determined from the caustic mass profile (see below) and σ_{200} is the projected velocity dispersion of galaxies within the caustics and within the projected radius r_{200} . Using this definition of contrast, there is no significant correlation of C_{200} with cluster mass for the CIRS clusters. Further, there is no significant difference between the distributions of the C_{200} statistic between the 400d-SDSS (excluding the four problematic groups) and CIRS samples (the samples differ at 59% confidence using a K-S test). This result suggests that any difference between the contrasts of the two samples is too subtle for our simple contrast statistic. We caution the reader that this contrast statistic is not well tested and may have subtle dependencies on the caustic method used to define membership. It

does provide a simple, quantitative way of comparing the 400d-SDSS sample to the CIRS sample.

We discuss the individual groups in more detail in §3.3.

3.2. Caustics and Mass Profiles

CAIRNS and CIRS showed that caustic masses of clusters agree well with mass estimates from both X-ray observations and Jeans’ analysis at small radii, where dynamical equilibrium is expected to hold (Rines et al. 2003, CIRS). Lokas & Mamon (2003) confirm that the mass of Coma estimated from higher moments of the velocity distribution agrees well with the caustic mass estimate (Geller et al. 1999). Recently, Diaferio et al. (2005) showed that caustic masses agree with weak lensing masses in three clusters at moderate redshift.

We briefly review the method DG and D99 developed to estimate the mass profile of a galaxy cluster by identifying caustics in redshift space. The method assumes




FIG. 3.— See Figure 2.

that clusters form in a hierarchical process. Application of the method requires only galaxy redshifts and sky coordinates. Toy models of simple spherical infall onto clusters produce sharp enhancements in the phase space density around the system. These enhancements, known as caustics, appear as a trumpet shape in scatter plots of redshift versus projected clustercentric radius (Kaiser 1987). DG and D99 show that random motions smooth out the sharp pattern expected from simple spherical infall into a dense envelope in the redshift-projected radius diagram (see also Vedel & Hartwick 1998). The edges of this envelope can be interpreted as the escape velocity as a function of radius. Galaxies outside the caustics are also outside the turnaround radius. The caustic technique provides a well-defined boundary between the infall region and interlopers; one may think of the technique as a method for defining membership that gives the cluster mass profile as a byproduct. In fact, the caustic technique can be used as a membership classification for any gravitationally bound system. For instance, Serra et al.

(2009) used caustics to identify interloper stars in five dwarf spheroidal galaxies, and Brown et al. (2009) use caustics to identify stars not bound to the Milky Way.

Operationally, we identify the caustics as curves which delineate a significant decrease in the phase space density of galaxies in the projected radius-redshift diagram. The details of the caustic technique used here are identical to those described in CIRS, with the following exception. Because 400d-SDSS groups are expected to be less massive than CIRS clusters, we isolate the groups initially by studying only galaxies within $R_p \leq 7h^{-1}\text{Mpc}$ and $\pm 4000\text{km s}^{-1}$ of the nominal group centers from the X-ray catalogs (compared to $10h^{-1}\text{Mpc}$ and $\pm 5000\text{km s}^{-1}$ for the CIRS clusters). We perform a hierarchical structure analysis to locate the centroid of the largest system in each volume. This analysis sometimes finds the center of another system in the field. In these cases, limiting the galaxies to a smaller radial and/or redshift range enables the algorithm to center on the desired group. For some groups, no satisfactory match between the hierarchical

TABLE 1
400D-SDSS BASIC PROPERTIES

400d ID	400d Name	X-ray Coordinates		$z_{\odot}$	$L_X/10^{43}$ erg s $^{-1}$	σ_p km s $^{-1}$	Flag	R_{comp} h^{-1} Mpc
		RA (J2000)	DEC (J2000)					
81	cl0810+4216	122.60085	42.27195	0.0640	1.128	473^{+73}_{-50}	2	>7
83 ^a	cl0820+5645	125.11083	56.75750	0.0430	0.049	636^{+79}_{-57}	1	0.2
89	cl0900+3920	135.01958	39.34000	0.0950	0.382	333^{+59}_{-39}	1	0.7
106	cl1010+5430	152.56584	54.50333	0.0462	0.052	266^{+53}_{-33}	1	2.1
112 ^a	cl1033+5703	158.46625	57.05306	0.0460	0.036	397^{+81}_{-51}	1	2.1
115	cl1039+3947	159.88042	39.79389	0.0930	0.202	193^{+132}_{-47}	0	0.3
118	cl1058+0136	164.55249	01.61583	0.0393	0.233	415^{+44}_{-33}	2	5.8
135	cl1159+5531	179.96333	55.53222	0.0810	0.575	321^{+79}_{-46}	1	0.2
155 ^a	cl1227+0858	186.80875	08.97083	0.0901	0.409	749^{+86}_{-64}	2	0.4
159	cl1236+1240	189.11917	12.40583	0.0440 ^b	0.054	384^{+117}_{-62}	0	4.0
170	cl1329+1143	202.36624	11.72306	0.0231	0.057	363^{+46}_{-33}	2	1.1
179 ^a	cl1343+5546	205.86624	55.78194	0.0685	0.112	465^{+54}_{-40}	2	0.3
199	cl1533+3108	233.32126	31.14861	0.0670	0.958	421^{+79}_{-51}	1	3.5
204	cl1630+2434	247.56125	24.58028	0.0637	0.823	465^{+74}_{-50}	2	>7
205	cl1631+2121	247.77042	21.36500	0.0952	0.331	518^{+171}_{-89}	0	2.0
227	cl2137+0026	324.27875	00.44667	0.0506	0.082	217^{+72}_{-38}	1	2.2

NOTE. — Col. (1) lists the number of the group in the 400d catalog (Burenin et al. 2007). Col. (2) lists the name of the group following the convention of Burenin et al. (2007). Col. (8) gives our subjective classification of the contrast between the group and foreground and background structure. A Flag of 2 indicates a “clean” infall pattern with little contaminating structure, while a Flag of 0 indicates a “weak” infall pattern for groups where the infall pattern is weakly present and/or significantly contaminated by nearby structure (see text for details). Col. (9) gives the projected radius of the nearest edge of the footprint of the SDSS spectroscopic survey.

^aStrongly affected by a nearby group or cluster. See §3.3 for details.

^bBurenin et al. (2007) list the redshift as $z=0.07$.

center and X-ray center is possible; for these groups we impose the X-ray center on the analysis (Table 2).

We discuss the groups individually in §3.3. Figures 2-3 show the caustics and Figures 4-5 show the associated mass profiles. Note that the caustics extend to different radii for different groups. D99 shows that the appearance of the caustics depends strongly on the line of sight; projection effects can therefore account for most of the differences in profile shape in Figures 2-3 without invoking non-homology among clusters. We use the caustics to determine group membership. Here, the term “group member” refers to galaxies both in the virial region and in the infall region and inside the caustics. Figures 2-3 show that the caustics effectively separate group members from background and foreground galaxies, although some interlopers may lie within the caustics. Note that, for some groups, the small samples of galaxies yield unrealistically low estimates of the uncertainties in the caustic mass profiles.

The D99 algorithm we use to identify the caustics generally agrees with the lines one would draw based on a visual impression. This consistency suggests that systematic uncertainties in the caustic technique are dominated by projection effects rather than the details of the algorithm. We now discuss the individual groups in more detail.

3.3. Comments on Individual Groups

Groups share many common features, but each system is unique. We describe the most relevant aspects of each 400d-SDSS group below.

- *cl0810+4216* The X-ray emission is centered on a $r=14.2$ ($M_r \approx -22.3$) galaxy. The hierarchical cen-

ter is located at the position of a strongly bound galaxy pair $387h^{-1}$ kpc to the NE. The X-ray center corresponds to the position of the Brightest Group Galaxy (BGG) and is probably the dynamical center of this system. The infall diagram produced when adopting the X-ray center as the system center suggests that four sub- L^* galaxies have slightly higher redshifts than the bulk of the group members. Excluding these galaxies from the caustic-selected sample would reduce the inferred velocity dispersion by $\sim 20\%$. The caustic technique classifies these four galaxies as members using either the X-ray or the hierarchical center.

- *cl0820+5645* The X-ray emission is centered on a pair of interacting galaxies with $r=15.7$ and $r=16.0$. A plot of group members on the sky reveals two main clumps of galaxies, one on the X-ray center and the other approximately centered on a $r = 14.6$ galaxy (MCG +09-14-024) without a redshift. This system is possibly a group-group merger.
- *cl0900+3920* The X-ray emission is centered $<1'$ from a pair of galaxies with $r=15.9$ and $r=17.1$. The Brightest Group Galaxy ($r=15.4$) lies $\sim 4'$ north of the X-ray center.
- *cl1010+5430* The X-ray emission is centered on a $r=14.2$ galaxy, UGC 06057. This galaxy has no SDSS redshift, but a redshift from Carballo et al. (1995) places it in the group.
- *cl1033+5703* The X-ray emission is centered on a $r=14.2$ galaxy. The spatial distribution of galax-




FIG. 4.— Comparison of caustic mass profiles to those estimates from the virial theorem and the projected mass estimator. The thick solid lines show the caustic mass profiles and the thin lines show the 1σ uncertainties in the mass profiles (for some systems, the thin lines blend with the thick lines). The axes are identical in all panels. The vertical bars indicate r_{200} and the maximum radius of the caustic mass profile (the smaller of r_{max} , the extent of the infall pattern, and r_t , the turnaround radius). Vertical dashed lines indicate r_t for groups where the infall pattern truncates before r_t . Red and green shaded regions show the formal 1σ uncertainties in the virial and projected mass profiles. Squares around group names indicate groups with contamination from nearby groups or clusters (see §3.3). Figure 5 shows similar plots for the rest of the sample.

ies near this group shows that the X-ray group is located $\sim 800h^{-1}\text{kpc}$ from a group of galaxies at the same redshift centered on CGCG 290-048 ($r=13.5$), a radio galaxy with a jet $1.3h^{-1}\text{Mpc}$ in extent (Masson 1979). The 400d ROSAT pointing does not include the CGCG 290-048 group, but it has been classified as an optically-selected group by Miller et al. (2005) and Merchán & Zandivarez (2005). The dynamics of cl1033+5703 are probably perturbed by the proximity of the CGCG 290-048 group (for an analogous configuration, see A2197E/W and A2199 Rines et al. 2002). Indeed, the CGCG 290-048 group displays a more symmetric infall pattern than does cl1033+5703, suggesting that the former system dominates the dynamics of the two systems.

- *cl1039+3947* The X-ray emission is approximately centered on a $r=15.0$ galaxy. A bright ($r=15.2$) spiral galaxy with several fainter companions lies about $2h^{-1}\text{Mpc}$ NW of the group center. This latter system may be an infalling group.
- *cl1058+0136* The X-ray emission is centered on UGC 06057, a $r=13.5$ galaxy without a SDSS redshift but confirmed as a group member using a redshift from Smith et al. (2000). This group is $6'6$ from the optical position of Abell 1139 (Abell et al. 1989) and previously has been identified with A1139 by Ebeling et al. (1998) and Böhringer et al. (2000b).
- *cl1159+5531* The X-ray emission is centered on a $r=14.1$ ($M_r \approx -23.0$) galaxy. This group was clas-




FIG. 5.— See Figure 4.

sified as an X-ray overluminous elliptical galaxy (OLEG) by Vikhlinin et al. (1999).

- *cl1227+0858* The X-ray emission is centered on a $r=15.0$ galaxy without a SDSS redshift. A redshift from Slingend et al. (1998) shows this galaxy to be a member. The group lies $\sim 0.7h^{-1}\text{Mpc}$ from the Brightest Cluster Galaxy (BCG) of Abell 1541, a cluster at the same redshift. The hierarchical center is coincident with the core of A1541, so we instead use the X-ray center to estimate the caustics. Böhringer et al. (2000b) report an X-ray flux of $(2.5 \pm 0.5) \times 10^{-12} \text{erg s}^{-1} \text{cm}^{-2}$ for A1541, ~ 6 times larger than the flux from cl1227+0858. This difference suggests that A1541 dominates the dynamics of cl1227+0858 (see cl1033+5703 above). The dynamical parameters reported for cl1227+0858 thus partially reflect the properties of A1541, a more massive and more X-ray luminous system. Consistent with this idea, 7 galaxies within $0.4h^{-1}\text{Mpc}$ of the X-ray center have a velocity dispersion

$(594^{+252}_{-111} \text{km s}^{-1})$ 20% smaller than the value reported in Table 1; at slightly larger radii there is a large spike in redshift space due to A1541 (Figure 3).

- *cl1236+1240* This extended X-ray source is approximately centered on IC 3574, a $r=14.1$ ($M_r \approx -22.4$) galaxy at $z=0.067$ (Burenin et al. 2007, no SDSS spectrum available). SDSS spectra, however, show that there are 15 galaxies at $z \approx 0.044$ within $0.5h^{-1}\text{Mpc}$ of the X-ray center, while there are no other galaxies within this radius at $z \approx 0.067$. The brightest of these galaxies is $r=15.5$ ($M_r \approx -20.2$). The X-ray emission is therefore associated with a single bright galaxy (similar to a fossil group, although most fossil groups do contain faint galaxies at the same redshift) or with a group of less luminous galaxies (or some combination). We adopt the latter interpretation that cl1236+1240 is associated with the group of less luminous galaxies.

TABLE 2
400D-SDSS HIERARCHICAL CENTERS AND OFFSETS FROM X-RAY CENTERS

Group	Hierarchical Center		Δcz^a	ΔR^a
	RA (J2000)	DEC (J2000)	km s ⁻¹	h ⁻¹ kpc
cl0810+4216	122.74532	42.33660	013	387
cl0820+5645 ^b	125.11101	56.75749	—	—
cl0900+3920	135.02003	39.34003	—	—
cl1010+5430	152.55563	54.55338	-366	115
cl1033+5703 ^b	158.46603	57.05308	—	—
cl1039+3947	159.87998	39.79392	—	—
cl1058+0136	164.53830	01.62267	-393	30
cl1159+5531	179.96301	55.53222	—	—
cl1227+0858 ^b	186.87500	08.82286	-34 ^c	— ^d
cl1236+1240	189.06404	12.44854	002	149
cl1329+1143	202.32590	11.65740	-17	89
cl1343+5546 ^b	205.90356	55.62635	-436	518
cl1533+3108	233.32102	31.14862	—	—
cl1630+2434	247.59420	24.57920	702	92
cl1631+2121	247.86281	21.51073	831	— ^d
cl2137+0026	324.25590	00.40342	127	121

^aNo value listed for the 6 groups where the X-ray center is used for the caustic analysis.

^bStrongly affected by a nearby group or cluster. See §3.3 for details.

^cBurenin et al. (2007) list a redshift of $z=0.087$ and reference Burke et al. (2003). However, Burke et al. (2003) list a redshift of $z=0.090$ for this group.

^dX-ray center adopted for spatial position and hierarchical central redshift. The hierarchical centers are offset on the sky by $\Delta R=685$ h⁻¹kpc and $753h^{-1}$ kpc for cl1227+0858 and cl1631+2121 respectively.

- *cl1329+1143* The X-ray emission is centered between NGC 5179 ($r=13.3$) and NGC 5171 ($r=12.9$). The latter has no SDSS spectrum, but a redshift from Falco et al. (1999) places NGC 5171 in the group. This group is also known as MKW 11 and its infall region was previously analyzed in CIRS. Note that the X-ray flux from the 400d catalog $[(1.3\pm0.2)\times10^{-12}\text{erg s}^{-1}\text{cm}^{-2}]$ is significantly smaller than the flux from RASS data Böhringer et al. (2000b) used in CIRS $[(5.7\pm0.8)\times10^{-12}\text{erg s}^{-1}\text{cm}^{-2}]$. The smaller flux from the 400d catalog would place this system below the flux limit used in CIRS. Osmond et al. (2004) analyze an XMM-Newton observation of this group and determine a flux of $(2.5\pm0.4)\times10^{-12}\text{erg s}^{-1}\text{cm}^{-2}$ in the (0.5–2.0 keV) band, intermediate between the RASS and 400d values. A possible explanation of the varying flux estimates is that a background group south-east of NGC 5171 (centered on a red galaxy with SDSS photometric redshift $z=0.21\pm0.01$) may have been included in the RASS flux, while the 400d flux may exclude some emission associated with group members. Osmond et al. (2004) further find evidence for interactions between the group members NGC 5171 and NGC 5176 and the surrounding gas. We find a slightly smaller velocity dispersion (363_{-33}^{+46} km s⁻¹) than do Osmond & Ponman (2004, 494 ± 99 km s⁻¹), who noted that their value of σ_p was larger than expected based on the measured $T_X=0.96\pm0.04$ keV. Our smaller value of σ_p mitigates this discrepancy. The NGC 5129 X-ray group lies $\sim3h^{-1}$ Mpc northeast of cl1329+1143 at a similar redshift; it is possible that these groups are physically associated.
- *cl1343+5546* The X-ray emission is centered on a $r=15.8$ galaxy. The group is located $11.4\approx500h^{-1}$ kpc from the optical center of A1783. The hierarchical center is coincident with the core of A1783, so the reported dynamical properties reflect those of the Abell cluster and not the 400d group. Unlike cl1227+0858/A1541 above, cl1343+5546 and A1783 are too close together to obtain a mass profile for cl1343+5546 alone. Ledlow et al. (2003) report an X-ray flux of $(0.51\pm0.12)\times10^{-12}\text{erg s}^{-1}\text{cm}^{-2}$ for A1783, ~2.5 times larger than the flux of cl1343+5546. The galaxies in A1783 exhibit significant substructure, perhaps indicating that the cluster is not yet relaxed.
- *cl1533+3108* The X-ray emission is centered on a $r=15.3$ galaxy possibly undergoing a merger with a $r=16.7$ galaxy; neither galaxy has a SDSS spectrum. A redshift from Small et al. (1997) places the brighter galaxy in the group. The central galaxy pair is surrounded by several bright galaxies. This group is also known as Abell 2092. Wu et al. (1999) list a velocity dispersion of $\sigma_p = 504_{-69}^{+115}$ km s⁻¹, compared with our estimate of $\sigma_p = 421_{-51}^{+79}$ km s⁻¹. Our estimate is based on a larger number of galaxies.
- *cl1630+2434* The X-ray emission is centered between a $r=14.6$ and a $r=15.2$ galaxy. The brighter galaxy (IC 4607) has no redshift available from either SDSS or the literature.
- *cl1631+2121* The X-ray center lies among seven red galaxies with $r=16.5$ –18.0 (one a spectroscop-

ically confirmed member); a $r=16.4$ spiral galaxy (and spectroscopically confirmed member) lies $\sim 1'$ northeast of the X-ray center. The hierarchical center is $\sim 750h^{-1}\text{kpc}$ northeast of the X-ray center, and $\sim 1'5$ southwest of a $r=15.4$ group member. Many group members lie close to the line connecting the X-ray and hierarchical centers. We center the mass profile on the X-ray center and the hierarchical redshift (Table 2). Additional redshifts, especially of the galaxies near the X-ray center, would help clarify the nature of this system.

- *cl2137+0026* The X-ray emission is centered between a $r=14.3$ early-type galaxy and a $r=14.4$ spiral galaxy. Both are group members, although the early-type galaxy has no SDSS spectrum (Falco et al. 1999).

3.4. Virial and Turnaround Masses and Radii

The caustic mass profiles allow direct estimates of the virial and turnaround radius in each group. For the virial radius, we estimate r_{200} (r_Δ is the radius within which the enclosed average mass density is $\Delta\rho_c$, where ρ_c is the critical density) by computing the enclosed density profile [$\rho(<r) = 3M(<r)/4\pi r^3$]; r_{200} is the radius which satisfies $\rho(<r_{200}) = 200\rho_c$. In our adopted cosmology, a system should be virialized inside the slightly larger radius $\sim r_{100} \approx 1.3r_{200}$ (Eke et al. 1996). We use r_{200} because it is more commonly used in the literature and thus allows easier comparison of results. For the turnaround radius r_t , we use equation (8) of Regös & Geller (1989) assuming $\Omega_m = 0.3$. For this value of Ω_m , the enclosed density is $3.5\rho_c$ at the turnaround radius. Varying Ω_m in the range 0.02–1 only changes the inferred value of r_t by $\pm 10\%$; the uncertainties in r_t from the uncertainties in the mass profile are comparable or larger (D99; Rines et al. 2002). If the w parameter in the equation of state of the dark energy ($P_\Lambda = w\rho_\Lambda$) satisfies $w \geq -1$, the dark energy has little effect on the turnaround overdensity (Gramann & Suhhonenko 2002).

Table 3 lists r_{200} , r_t , and the masses M_{200} and M_t enclosed within these radii. For cl0820+0945, cl1329+1143, and cl1631+21214, the maximum extent of the caustics r_{max} is smaller than r_t . For these groups, r_t is a minimum value assuming that there is no additional mass outside r_{max} . The best estimate of the mass contained in infall regions clearly comes from those groups for which $r_{max} \geq r_t$. The average mass within the turnaround radius for these groups is 2.4 ± 0.4 times the virial mass M_{200} , demonstrating that groups are still forming in the present epoch and consistent with the CIRS estimate of 2.19 ± 0.18 . Similarly, the average turnaround radius is $(5.01 \pm 0.25)r_{200}$ for groups with $r_{max} \geq r_t$, again in agreement with the CIRS result of $(4.96 \pm 0.08)r_{200}$. Simulations of the future growth of large-scale structure (Gramann & Suhhonenko 2002; Nagamine & Loeb 2003; Busha et al. 2003) for our assumed cosmology ($\Omega_m = 0.3, \Omega_\Lambda = 0.7$) suggest that galaxies currently inside the turnaround radius of a system will continue to be bound to that system. Our results for the turnaround radius and mass agree with the predictions of Busha et al. (2005), who find that the ultimate mass of dark matter haloes in simulations is $1.9M_{200}$. The agreement between the estimates of M_t/M_{200} and r_t/r_{200} between the

CIRS and 400d-SDSS samples suggests that the overall shapes of cluster mass profiles into their infall regions are not strongly dependent on cluster mass into the group regime. We compare the virial masses M_{200} from the caustics with masses calculated using the virial theorem in §3.5.

One striking result of this analysis is that the caustic pattern is often visible beyond the turnaround radius of a group (similar to some CIRS clusters). This result suggests that even groups may have strong dynamic effects on surrounding large-scale structure beyond the turnaround radius. For our assumed cosmology, this large-scale structure is probably not bound to the group.

3.5. Comparison to Virial and Projected Mass Estimates

We apply the virial mass and projected mass estimators (Heisler et al. 1985) to the 400d-SDSS groups. For the latter, we assume the galaxies are on isotropic orbits. We must define a radius of virialization within which the galaxies are relaxed. We use r_{200} (Table 3) and include only galaxies within the caustics. We thus assume that the caustics provide a good division between group galaxies and interlopers (see Figures 2-3). For details on these mass calculations, see CIRS. Table 4 lists the virial and projected mass estimates.

Figure 6 compares the virial and caustic mass estimates at r_{200} . The mean ratios of these estimates are $M_v/M_c = 0.91 \pm 0.12$. The caustic mass estimates are consistent with virial mass estimates even assuming a correction factor $C \approx 0.1 - 0.2M_{vir}$ for the surface pressure, consistent with the best-fit NFW profiles (see also Carlberg et al. 1997; Girardi et al. 1998; Koranyi & Geller 2000; Rines et al. 2003, CIRS). Note that Lemze et al. (2009) suggest a possible modification of the form factor $\mathcal{F}_\beta$ used in the caustic technique based on a comparison of the caustic and lensing mass profiles of A1689. Lopes et al. (2009) find a smaller scatter in scaling relations using virial masses rather than caustic masses for the CIRS clusters.

Figures 4-5 compare the mass profiles estimated from the caustics, virial theorem, and projected mass estimator. The virial and projected mass profiles $M(<R_P)$ are calculated using all galaxies within the projected radius R_P (note that the virial assumptions hold only at the virial radius). The projected mass estimator consistently overestimates the mass at small radii and underestimates the mass at large radii relative to the other profiles. This behavior suggests that this estimator is best for estimating virial masses but not mass profiles. The virial and caustic mass profiles generally agree although there are many groups with large disagreements. The caustic mass profiles do not appear to consistently overestimate or underestimate the mass relative to the virial mass profiles. This result supports our use of caustic mass profiles as a tracer of the total group mass profile in the next section.

3.6. The Shapes of Group Mass Profiles

We fit the mass profiles of the 400d-SDSS groups to three simple analytic models. The simplest model of a self-gravitating system is a singular isothermal sphere (SIS). The mass of the SIS increases linearly with radius. Navarro et al. (1997) and Hernquist (1990) propose two-parameter models based on CDM simulations of haloes.

TABLE 3
400D-SDSS CHARACTERISTIC RADII AND MASSES

Group	r_{500} $h^{-1}\text{Mpc}$	r_{200} $h^{-1}\text{Mpc}$	r_t $h^{-1}\text{Mpc}$	r_{max} $h^{-1}\text{Mpc}$	M_{200} $10^{14}M_{\odot}$	M_t $10^{14}M_{\odot}$	M_t/M_{200}
cl0810+4216	0.52	0.83	4.05	9.49	1.32 ± 0.05	2.70 ± 0.14	2.05
cl0820+5645 ^a	0.67	1.09	>4.55	3.94	3.03 ± 0.02	3.84 ± 0.02	1.27
cl0900+3920	0.35	0.58	4.17	9.19	0.45 ± 0.06	2.95 ± 0.48	6.58
cl1010+5430	0.20	0.42	2.66	9.80	0.17 ± 0.04	0.77 ± 0.25	4.45
cl1033+5703 ^a	0.38	0.60	3.14	9.60	0.51 ± 0.02	1.26 ± 0.06	2.48
cl1039+3947	0.29	0.45	2.08	3.64	0.21 ± 0.04	0.37 ± 0.08	1.74
cl1058+0136	0.51	0.86	4.17	>10	1.51 ± 0.00	2.96 ± 0.00	1.96
cl1159+5531	0.39	0.63	2.86	9.29	0.57 ± 0.00	0.95 ± 0.00	1.67
cl1227+0858 ^a	0.56	0.97	4.09	4.34	2.13 ± 0.83	2.79 ± 1.16	1.31
cl1236+1240	0.49	0.71	2.88	9.70	0.84 ± 0.00	0.97 ± 0.00	1.16
cl1329+1143	0.40	0.59	>2.66	1.82	0.48 ± 0.00	0.77 ± 0.00	1.58
cl1343+5546 ^a	0.53	0.90	4.70	>10	1.69 ± 0.06	4.23 ± 0.20	2.50
cl1533+3108	0.32	0.54	2.50	2.63	0.36 ± 0.23	0.64 ± 0.47	1.76
cl1630+2434	0.52	0.76	3.12	9.09	1.03 ± 0.18	1.24 ± 0.23	1.20
cl1631+2121	0.40	0.60	>2.72	1.82	0.49 ± 0.39	0.82 ± 0.71	1.67
cl2137+0026	0.35	0.51	2.67	8.99	0.32 ± 0.01	0.78 ± 0.05	2.45

NOTE. — Cols. (1)-(3) list r_{δ} for three values of overdensity δ , where r_{δ} is the radius within which the enclosed mass density is δ times the critical density (we adopt $\delta=3.5$ as the turnaround overdensity). Col. (4) gives the turnaround radius r_t for each group. Col. (5) gives the maximum radius r_{max} within which the caustics are detected. Note that when $r_{max} < r_t$, the estimate of r_t is an underestimate if any mass lies between r_{max} and r_t .

^aStrongly affected by a nearby group or cluster. See §3.3 for details.

TABLE 4
400D-SDSS VIRIAL AND PROJECTED MASSES

Group	r_{200} $h^{-1}\text{Mpc}$	M_{200} $10^{14}M_{\odot}$	M_{proj} $10^{14}M_{\odot}$	M_{vir} $10^{14}M_{\odot}$
cl0810+4216	0.83	1.32 ± 0.05	0.79 ± 0.19	1.14 ± 0.17
cl0820+5645 ^a	1.09	3.03 ± 0.02	3.15 ± 0.65	3.15 ± 0.41
cl0900+3920	0.58	0.45 ± 0.06	0.21 ± 0.06	0.27 ± 0.04
cl1010+5430	0.42	0.17 ± 0.04	0.20 ± 0.06	0.12 ± 0.02
cl1033+5703 ^a	0.60	0.51 ± 0.02	0.45 ± 0.14	0.35 ± 0.06
cl1039+3947	0.45	0.21 ± 0.04	0.08 ± 0.04	0.18 ± 0.00
cl1058+0136	0.86	1.51 ± 0.00	1.02 ± 0.18	0.76 ± 0.09
cl1159+5531	0.63	0.57 ± 0.00	0.24 ± 0.08	0.41 ± 0.08
cl1227+0858 ^a	0.97	2.13 ± 0.83	1.09 ± 0.21	3.58 ± 0.45
cl1236+1240	0.71	0.84 ± 0.00	0.39 ± 0.16	0.59 ± 0.12
cl1329+1143	0.59	0.48 ± 0.00	0.58 ± 0.12	0.52 ± 0.07
cl1343+5546 ^a	0.90	1.69 ± 0.06	0.79 ± 0.15	0.85 ± 0.11
cl1533+3108	0.54	0.36 ± 0.23	0.46 ± 0.13	0.63 ± 0.10
cl1630+2434	0.76	1.03 ± 0.18	0.35 ± 0.09	0.64 ± 0.10
cl1631+2121	0.60	0.49 ± 0.39	0.31 ± 0.13	0.97 ± 0.20
cl2137+0026	0.51	0.32 ± 0.01	0.07 ± 0.03	0.09 ± 0.02

NOTE. — Cols. (4) and (5) give the projected and virial mass estimates for each group computed with all galaxies inside the radius r_{200} determined from the caustic mass profile.

^aStrongly affected by a nearby group or cluster. See §3.3 for details.

We note that the caustic mass profiles mostly sample large radii and are therefore not very sensitive to the inner slope of the mass profile. Thus, we do not consider alternative models which differ only in the inner slope of the density profile (e.g., Moore et al. 1999). At large radii, the best constraints on cluster mass profiles come from galaxy dynamics and weak lensing. The caustic mass profiles of Coma (Geller et al. 1999), A576 (Rines et al. 2000), A2199 (Rines et al. 2002) and the rest of the CAIRNS and CIRS clusters (Rines et al. 2003; Rines & Diaferio 2006) provided strong evidence against a singular isothermal sphere (SIS) profile and in favor of steeper mass density profiles predicted by Navarro et al. (1997, NFW) and Hernquist (1990). Only recently have weak lensing mass estimates been able to distinguish between

SIS and NFW density profiles at large radii (Clowe & Schneider 2001; Kneib et al. 2003; Mandelbaum et al. 2006).

At large radii, the NFW mass profile increases as $\ln(r)$ and the mass of the Hernquist model converges. The NFW mass profile is

$$M(<r) = \frac{M(a)}{\ln(2) - \frac{1}{2}} \left[\ln\left(1 + \frac{r}{a}\right) - \frac{r}{a+r} \right] \quad (1)$$

where a is the scale radius and $M(a)$ is the mass within a . We fit the parameter $M(a)$ rather than the characteristic density δ_c ($M(a) = 4\pi\delta_c\rho_c a^3 [\ln(2) - \frac{1}{2}]$ where ρ_c is the critical density) because $M(a)$ and a are much less correlated than δ_c and a (Mahdavi et al. 1999). The




FIG. 6.— Caustic masses at r_{200} compared to virial masses at the same radius. Symbol types are the same as for Figure 1. Four groups that are strongly affected by nearby systems are highlighted with large circles. Errorbars show 1σ uncertainties and the solid line shows a ratio of unity.

Hernquist mass profile is

$$M(< r) = M \frac{r^2}{(r + a_H)^2} \quad (2)$$

where a_H is the scale radius and M is the total mass. Note that $M(a_H) = M/4$. The SIS mass profile is $M(< r) \propto r$. We minimize χ^2 and list the best-fit parameters a , r_{200} , the concentration $c_{200}=r_{200}/a$, and M_{200} for the best-fit NFW model and indicate the best-fit profile type in Table 5. We also list the parameter $c_{101}=r_{101}/a$; some authors prefer to use r_{101} as the virial radius for this cosmology. We perform the fits on all data points within the maximum radial extent of the caustics r_{max} listed in Table 3 and with caustic amplitude $\mathcal{A}(r) > 100 \text{ km s}^{-1}$.

Because the individual points in the mass profile are not independent, the absolute values of χ^2 are indicative only, but it is clear that the NFW and Hernquist profiles provide acceptable fits to the caustic mass profiles; the SIS is never the best-fit profile and it is actually excluded for nearly all groups. The NFW profile provides a better fit to the data than the Hernquist profile for 5 of the 16 400d-SDSS groups; the remaining 11 are better fit by a Hernquist profile.

Figure 7 shows the shapes of the caustic mass profiles scaled by r_{200} and M_{200} along with SIS, NFW, and Hernquist model profiles. The colored lines show different model mass profiles. The straight dashed line is the SIS, the solid lines are NFW profiles with $c_{200}=3, 5$, and 10, and the curved dashed lines are Hernquist profiles with two different scale radii. The average value of c_{200} is 6.5, consistent with CIRS and with the values expected from simulations for clusters (NFW; Bullock et al. 2001). All three model profiles agree fairly well with the caustic mass profiles in the range $(0.1-1)r_{200}$. The SIS only fails beyond $\sim 1.5r_{200}$; this is why lensing has had trouble distinguishing between SIS and NFW profiles. As discussed in D99, the caustic technique can be subject to large variations for individual groups due to projection effects. The best constraints on the shapes of cluster

and group mass profiles are obtained by averaging over many lines of sight, or for real observations, over many different clusters and groups.

The concentration parameters $c_{200} = r_{200}/a$ for the NFW models are in the range 1-15 (Table 5), in good agreement with the predictions of numerical simulations (Navarro et al. 1997). Numerical simulations suggest that c should decrease slightly ($\sim 30\%$) with increasing mass over this mass range, but this decrease is comparable to the scatter in c in simulated clusters (Navarro et al. 1997; Bullock et al. 2001). The errorbars in Figure 8 indicate the average values and 1σ scatter in $c_{101} = r_{101}/a$ in simulations (Bullock et al. 2001).

Observations of mass profiles using X-ray mass estimates (Buote et al. 2007) and weak lensing (Comerford & Natarajan 2007) appear to confirm a decrease in c with increasing mass, although some massive clusters apparently have high concentrations (e.g. Lemze et al. 2009). In contrast, the CIRS clusters showed a weak positive correlation of c with mass (Figure 8). The 400d-SDSS groups do not confirm this correlation (Figure 8). A Spearman rank-sum test indicates that there is no significant correlation between c_{101} and M_{101} (20% confidence) for the 400d-SDSS groups. A Spearman test for the CIRS clusters suggests a correlation at the 92.7% confidence level. By combining the CIRS and 400d-SDSS data, a Spearman test indicates a correlation between concentration and mass at the 93.6% confidence level (correlation coefficient 0.19). This result indicates possible tension between caustic mass profiles and simulations, as the caustic mass profiles indicate that concentrations increase rather than decrease with increasing cluster mass. More data for groups at lower masses would help clarify the situation.

To further explore the dependence of mass profile shapes on mass, the right panel of Figure 7 shows the scaled mass profiles of the most massive quartile of CIRS clusters. Comparing the two panels reveals no obvious trend with increasing mass: the profiles of 400d-SDSS groups have larger scatter, but the average of the profiles appears similar to that of the massive CIRS clusters.

4. SCALING RELATIONS OF CLUSTERS AND GROUPS

Scaling relations between simple cluster/group observables and masses provide insight into the nature of cluster/group assembly and the properties of various cluster/group components. Establishing these relations for local clusters and groups is critical for future studies of clusters in the distant universe with the goal of constraining dark energy (Majumdar & Mohr 2004; Lin et al. 2004).

We apply the prescription of Danese et al. (1980) to determine the mean redshift $cz_\odot$ and projected velocity dispersion σ_p of each group from all galaxies within the caustics. We calculate σ_p using only the group members projected within r_{200} estimated from the caustic mass profile. Note that our estimates of r_{200} do not depend directly on σ_p .

One of the simplest X-ray observables of clusters and groups is X-ray luminosity. For optical data, the projected velocity dispersion σ_p is a straightforward observable. In particular, Lopes et al. (2009) reanalyze the CIRS clusters and find that the velocity dispersions are

TABLE 5
400d-SDSS MASS PROFILE FIT PARAMETERS

Group	a_{NFW} $h^{-1}\text{Mpc}$	r_{200} $h^{-1}\text{Mpc}$	c_{200}	M_{200} $10^{14} M_{\odot}$	Best-fit Profile	c_{101}
cl0810+4216	0.185	0.79	4.26	1.14	N	5.75
cl0820+5645 ^a	0.205	0.95	4.63	1.98	H	6.23
cl0900+3920	0.590	0.60	1.02	0.50	N	1.49
cl1010+5430	0.415	0.40	0.96	0.15	H	1.42
cl1033+5703 ^a	0.160	0.61	3.83	0.53	H	5.19
cl1039+3947	0.072	0.41	5.75	0.16	H	7.68
cl1058+0136	0.297	0.77	2.59	1.06	H	3.58
cl1159+5531	0.160	0.58	3.64	0.46	H	4.95
cl1227+0858 ^a	0.415	0.75	1.80	1.01	H	2.59
cl1236+1240	0.081	0.62	7.70	0.57	H	10.20
cl1329+1143	0.064	0.60	9.32	0.49	N	12.30
cl1343+5546 ^a	0.200	0.86	4.29	1.46	N	5.78
cl1533+3108	0.138	0.48	3.46	0.25	H	4.71
cl1630+2434	0.051	0.68	13.40	0.75	H	17.56
cl1631+2121	0.092	0.58	6.32	0.47	H	8.52
cl2137+0026	0.081	0.50	6.20	0.30	N	8.27

NOTE. — Cols. (2) and (3) give the scale radius a_{NFW} and r_{200} of the best-fit NFW profile. Cols. (4) and (6) give the NFW concentration parameters $c_{\delta} = r_{\delta}/a_{NFW}$ where r_{δ} is the radius within which the enclosed density is δ times the critical density.

^aStrongly affected by a nearby group or cluster. See §3.3 for details.




FIG. 7.— Scaled caustic mass profiles for the 400d-SDSS groups (excluding those contaminated by nearby large-scale structure) compared to simple models. The thin solid lines show the caustic mass profiles normalized by r_{200} and M_{200} . The long-dashed line shows a singular isothermal sphere, the colored solid lines show NFW profiles (with concentrations $c_{200}=3,5,10$ from top to bottom at large radii). The short-dashed lines are Hernquist profiles with scale radii different by a factor of two. Right: Scaled caustic mass profiles for the most massive quartile (ordered by M_{200}) of CIRS clusters with the same model profiles.

robust to the particular choice of interloper removal algorithms, while their virial masses have significant scatter relative to the CIRS values (primarily due to differences in estimates of r_{200}). The $L_X - \sigma_p$ relation is expected to have the form $L_X \propto \sigma_p^4$ under simple theoretical models (Quintana & Melnick 1982). For clusters, most recent estimates of the slope of this relation are consistent with a slope of 4. For galaxy groups, however, some authors find shallower slopes (Mahdavi et al. 2000; Xue & Wu 2000), while others find slopes comparable to clusters (Ponman et al. 1996; Mulchaey & Zabludoff 1998). In the Group Evolution Multiwavelength Survey (GEMS), Osmond & Ponman (2004) find a slope of 2.31 ± 0.61 for groups, but the slope for a combined sample of groups and clusters

is 4.55 ± 0.25 . In a similar vein, Mahdavi et al. (2000) find that the best-fit relation for a sample of groups and clusters is a broken power-law with a shallower slope for groups than for clusters. Unusual systems may affect the $L_X - \sigma_p$ relation: for instance, Helsdon et al. (2005) show that galaxy groups with apparently very small σ_p and large L_X may have larger σ_p than earlier estimates, bringing these groups more in line with a steeper relation. Khosroshahi et al. (2007) find that fossil groups follow a $L_X - \sigma_p$ relation with slope 2.74 ± 0.45 , shallower than clusters (they also note a tendency for fossil groups to have slightly larger L_X for a given σ_p than non-fossil groups). In the 400d-SDSS sample, we selected the groups based on their X-ray emission, whereas most pre-




FIG. 8.— Concentrations $c_{101}=r_{101}/a$ of the best-fit NFW profiles versus M_{101} for the 400d-SDSS groups (solid hexagons) and the CIRS clusters (open circles). The dashed lines show the expected trend from (thick dashed line) simulations and a parametric model (thin dashed line Bullock et al. 2001). The solid and dash-dotted lines show the median value of c_{101} and $\log(c_{101})$ for the CIRS clusters as a function of cluster mass.




FIG. 9.— Velocity dispersions at r_{200} compared to X-ray luminosities. The solid line is the bisector of the least squares fits for the CIRS clusters. The dashed lines show the $\sigma_p - L_X$ relations from RASS-SDSS (Popesso et al. 2005). Symbol types are explained in Figure 1; large open circles highlight groups strongly affected by nearby structures. The two arrows indicate the locations of these data points for the larger clusters located close to two of these groups (A1541 near cl1227+0858 and A1783 near cl1343+5546).

vious studies (Ponman et al. 1996; Mulchaey & Zabludoff 1998; Mahdavi et al. 2000; Osmond & Ponman 2004) begin with catalogs of optically selected groups and then analyze either pointed X-ray observations or RASS data.

The 400d-SDSS X-ray luminosities are measured in the ROSAT band (0.1-2.4 keV) and corrected for Galactic absorption (see Burenin et al. 2007, for details). Figure 9 shows the $L_X - \sigma_p$ relation for the CIRS clusters along with the best-fit $L_X - \sigma_p$ relation of the RASS-SDSS (Popesso et al. 2005) and CIRS clusters, the GEMS

groups, and the GEMS groups+clusters fit (Osmond & Ponman 2004). The 400d-SDSS groups roughly follow the relations of the RASS-SDSS and CIRS samples, but the scatter is large. Squares, triangles, and crosses in Figure 9 show systems from CIRS, Mahdavi et al. (2000), and Osmond & Ponman (2004) respectively. Note that the 400d observations are typically not deep enough to separate group emission from contributions from individual galaxy halos, so the X-ray luminosities are not necessarily directly comparable to those in some other studies (e.g., Mahdavi et al. 2000; Osmond & Ponman 2004). However, because the 400d-SDSS groups lie close to RASSCALs and GEMS groups, this effect is probably small.

The groups previously identified as undergoing mergers are highlighted with large circles in Figure 9. These groups are the most extreme outliers from the CIRS and RASS-SDSS scaling relations and the locus of points from earlier studies. This result can be easily understood as a result of confusing an L_X measured for a sub-cluster with the velocity dispersion σ_p reflecting the velocity dispersion of the parent cluster. Consistent with this explanation, the arrows in Figure 9 show where cl1227+0858 (cl1343+5546) would appear if we used the X-ray luminosity of the nearby, more luminous system A1541 (A1783); both points move closer to the CIRS $L_X - \sigma$ relation. Note that only one of the 400d-SDSS groups (cl1159+5531) is a likely fossil group (Vikhlinin et al. 1998), so the sample is not dominated by these unusual systems. Specifically, Voevodkin et al. (2009) search for fossil groups and clusters in the 400d survey; cl1159+5531 is the only fossil group at $z < 0.1$, and it only qualifies under a relaxed definition of fossil groups.

Excluding the four groups contaminated by large-scale structure, the bisector of the least-squares fits has a slope of 3.4 ± 1.6 , intermediate between the shallow slopes found by some investigators for groups and the steeper slopes found for clusters (see Figure 9). Because the uncertainty in the slope for the 400d-SDSS groups is large, the slope is not significantly different from any of these relations.

We test the consistency of the $L_X - \sigma_p$ relation for 400d-SDSS groups with that of the CIRS clusters by fitting these relations for the CIRS clusters alone and for the CIRS clusters with the 400d-SDSS groups (excluding those contaminated by large-scale structure) added to the sample. For the CIRS data alone, the slope of the bisector of least-squares fits is 3.7 ± 0.6 , and the slope for the combined CIRS and 400d-SDSS data is 4.1 ± 0.4 .

From the limited sample, we conclude that the X-ray-selected 400d-SDSS groups have an $L_X - \sigma_p$ relation consistent with previous studies of both larger-mass clusters (e.g., CIRS) and smaller-mass groups (e.g., GEMS). We find no evidence of large biases in this relation between X-ray-selected and optically-selected systems. An important caveat to this conclusion is that some recent studies of clusters (Popesso et al. 2007) and groups (Rasmussen et al. 2006) find that there may be some systems that are significantly “X-ray underluminous,” perhaps because these systems are dynamically young.

Figure 10 shows L_X versus M_{500} as estimated from the caustics for the 400d-SDSS groups. Figure 10 also shows the data from CIRS and the best-fit $M_{500} - L_X$ relations for CIRS (caustic masses) and RASS-SDSS (both virial




FIG. 10.— Left: Caustic masses at r_{500} compared to X-ray luminosities. The solid line is the bisector of the ordinary least squares fits to the CIRS data. The red and blue lines show the $M_{500} - L_X$ relations for RASS-SDSS (Popesso et al. 2005) for optical and X-ray masses respectively. Symbol types are explained in Figure 1; large open circles highlight groups strongly affected by nearby structures. The two arrows indicate the locations of these data points for the larger clusters located close to two of these groups (A1541 near cl1227+0858 and A1783 near cl1343+5546). Right: Same for $M_{200} - L_X$.

masses and T_X -based masses Popesso et al. 2005).

Figure 11 shows the $M_{200} - \sigma_p$ relation. The tight relation indicates that the caustic masses are well correlated with velocity dispersion estimates. The good correlation is perhaps not surprising because both parameters depend on the galaxy velocity distribution. The best-fit slope for the CIRS clusters is $M_{200} \propto \sigma_p^{3.18 \pm 0.19}$ with the uncertainty estimated from jackknife resampling. The dashed line in Figure 11 shows the virial mass- σ_p relation for dark matter halos in simulations (Evrard et al. 2008, slope 3.035 ± 0.023). The agreement between these relations from CIRS and simulations further supports the use of caustics as a mass estimator (we compare the caustic masses to virial mass estimates in §3.5). The four merging groups in 400d-SDSS represent the most extreme outliers from the CIRS relation. The remaining 400d-SDSS groups approximately follow the CIRS relations, but we do not attempt to fit a relation for the 400d-SDSS groups.

In the previous section we found that many of the 400d groups are either group-group mergers (cl0820+5645) or group-cluster mergers (cl1033+5703, cl1227+0858, cl1343+5546). These groups represent the most extreme outliers in the $L_X - \sigma_p$ relation (Figure 9) and the $L_X - M$ relations (Figure 10). This result reinforces the idea that a detailed understanding of the large-scale structure surrounding low-mass systems is critical to obtaining accurate scaling relations. Previous studies, (e.g., Helsdon & Ponman 2000; Mahdavi et al. 2000) usually exclude such systems for similar reasons.

5. DISCUSSION

We use the Fifth Data Release of the Sloan Digital Sky Survey to study galaxy groups and compare their properties to more massive clusters. In particular, we study the presence of infall patterns around galaxy groups, use these patterns to measure mass profiles, and use these masses to compare the scaling relations of groups and clusters. This study extends our previous work on infall




FIG. 11.— Caustic masses at r_{200} compared to velocity dispersions within r_{200} . The solid line is the bisector of the ordinary least squares fits for the CIRS clusters. Symbol types are explained in Figure 1; large open circles highlight groups strongly affected by nearby structures. The dashed line shows the virial mass- σ_p relation for dark matter halos in simulations (Evrard et al. 2008).

patterns around X-ray clusters in SDSS (Rines & Diaferio 2006) to smaller X-ray fluxes more typical of groups. These systems typically have smaller X-ray luminosities and masses than the systems studied in previous investigations. Despite their smaller masses (compared to CAIRNS and CIRS clusters), well-defined infall patterns are present in most of these groups. Four of the 16 groups are contaminated by large-scale structure, preventing an accurate determination of their infall patterns.

We use the infall patterns to compute mass profiles for the groups and compare them to model profiles. Group infall regions are well fit by NFW and Hernquist profiles

and poorly fit by singular isothermal spheres, similar to the results from CAIRNS and CIRS for clusters. Observed clusters and groups resemble those in simulations, and their mass profiles are well described by extrapolations of NFW or Hernquist models out to the turnaround radius. The shapes of the best-fit NFW group mass profiles agree reasonably well with the predictions of simulations; the average concentration is $c_{200} \approx 6.5$, slightly smaller than the value for group size halos extrapolated from Bullock et al. (2001) and with similar scatter. These mass profiles test the shapes of dark matter haloes on a scale difficult to probe with weak lensing or any other mass estimator.

There has been much discussion of whether galaxy groups follow different scaling relations than clusters, particularly with respect to the $L_X - \sigma_p$ relation (e.g., Mahdavi et al. 2000; Helsdon & Ponman 2000; Osmond & Ponman 2004). Using the caustics to determine group membership, velocity dispersions, and mass profiles, the 400d-SDSS groups generally follow the same scaling relations as clusters, but the small number of 400d-SDSS groups does not allow us to exclude results suggesting that groups have a shallower $L_X - \sigma_p$ relation.

Understanding scaling relations of clusters and groups is critical for future attempts to constrain dark energy with cluster abundance measurements. Detailed mass estimates are difficult to obtain for groups and low-mass clusters because their X-ray emission is typically faint and because they contain fewer galaxies than massive clusters. This work represents part of an effort to increase the sample of groups and low-mass clusters with detailed mass estimates. Future work is needed to measure cluster scaling relations robustly across a wide mass range. In particular, deep high-resolution X-ray obser-

vations of groups like those studied here could enable measurements of group masses and X-ray temperatures, while additional optical spectroscopy could improve measurements of velocity dispersions and dynamical masses.

We thank the referee for a careful reading of the manuscript and for several helpful suggestions that significantly improved the presentation of the results. We thank Margaret Geller for useful discussions during early stages of this work and for insightful comments on early drafts. Support from the PRIN2006 grant “Costituenti fondamentali dell’Universo” of the Italian Ministry of University and Scientific Research and from the INFN grant PD51 is gratefully acknowledged. Funding for the Sloan Digital Sky Survey (SDSS) has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Aeronautics and Space Administration, the National Science Foundation, the U.S. Department of Energy, the Japanese Monbukagakusho, and the Max Planck Society. The SDSS Web site is <http://www.sdss.org/>. The SDSS is managed by the Astrophysical Research Consortium (ARC) for the Participating Institutions. The Participating Institutions are The University of Chicago, Fermilab, the Institute for Advanced Study, the Japan Participation Group, The Johns Hopkins University, the Korean Scientist Group, Los Alamos National Laboratory, the Max-Planck-Institute for Astronomy (MPIA), the Max-Planck-Institute for Astrophysics (MPA), New Mexico State University, University of Pittsburgh, University of Portsmouth, Princeton University, the United States Naval Observatory, and the University of Washington.

REFERENCES

- Abell, G. O., Corwin, H. G., & Olowin, R. P. 1989, *ApJS*, 70, 1
- Adelman-McCarthy, J. et al. 2007, *ApJS*, 172, 634
- Böhringer, H. et al. 2000a, *ApJS*, 129, 435
- . 2000b, *ApJS*, 129, 435
- Biviano, A. & Girardi, M. 2003, *ApJ*, 585, 205
- Brown, W. R., Geller, M. J., Kenyon, S. J., & Diaferio, A. 2009, submitted to *AJ*
- Bullock, J. S. et al. 2001, *MNRAS*, 321, 559
- Buote, D. A., Gastaldello, F., Humphrey, P. J., Zappacosta, L., Bullock, J. S., Brighenti, F., & Mathews, W. G. 2007, *ApJ*, 664, 123
- Burenin, R. A., Vikhlinin, A., Hornstrup, A., Ebeling, H., Quintana, H., & Mescheryakov, A. 2007, *ApJS*, 172, 561
- Burke, D. J., Collins, C. A., Sharples, R. M., Romer, A. K., & Nichol, R. C. 2003, *MNRAS*, 341, 1093
- Busha, M. T., Adams, F. C., Wechsler, R. H., & Evrard, A. E. 2003, *ApJ*, 596, 713
- Busha, M. T., Evrard, A. E., Adams, F. C., & Wechsler, R. H. 2005, *MNRAS*, 363, L11
- Carballo, R., Warwick, R. S., Barcons, X., Gonzalez-Serrano, J. I., Barber, C. R., Martinez-Gonzalez, E., Perez-Fournon, I., & Burgos, J. 1995, *MNRAS*, 277, 1312
- Carlberg, R. G., Yee, H. K. C., & Ellingson, E. 1997, *ApJ*, 478, 462
- Clowe, D. & Schneider, P. 2001, *A&A*, 379, 384
- Comerford, J. M. & Natarajan, P. 2007, *MNRAS*, 379, 190
- Cross, N. J. G., Driver, S. P., Liske, J., Lemon, D. J., Peacock, J. A., Cole, S., Norberg, P., & Sutherland, W. J. 2004, *MNRAS*, 349, 576
- Dai, X., Kochanek, C. S., & Morgan, N. D. 2007, *ApJ*, 658, 917
- Danese, L., de Zotti, G., & di Tullio, G. 1980, *A&A*, 82, 322
- Diaferio, A. 1999, *MNRAS*, 309, 610
- . 2009, *ArXiv e-prints*
- Diaferio, A. & Geller, M. J. 1997, *ApJ*, 481, 633
- Diaferio, A., Geller, M. J., & Rines, K. J. 2005, *ApJ*, 628, L97
- Drinkwater, M. J., Gregg, M. D., & Colless, M. 2001, *ApJ*, 548, L139
- Ebeling, H., Edge, A. C., Böhringer, H., Allen, S. W., Crawford, C. S., Fabian, A. C., Voges, W., & Huchra, J. P. 1998, *MNRAS*, 301, 881
- Eke, V. R., Cole, S., & Frenk, C. S. 1996, *MNRAS*, 282, 263
- Evrard, A. E. et al. 2008, *ApJ*, 672, 122
- Falco, E. E. et al. 1999, *PASP*, 111, 438
- Geller, M. J., Diaferio, A., & Kurtz, M. J. 1999, *ApJ*, 517, L23
- Girardi, M., Giuricin, G., Mardirossian, F., Mezzetti, M., & Boschin, W. 1998, *ApJ*, 505, 74
- Gramann, M. & Suhhonenko, I. 2002, *MNRAS*, 337, 1417
- Heisler, J., Tremaine, S., & Bahcall, J. N. 1985, *ApJ*, 298, 8
- Helsdon, S. F. & Ponman, T. J. 2000, *MNRAS*, 319, 933
- Helsdon, S. F., Ponman, T. J., & Mulchaey, J. S. 2005, *ApJ*, 618, 679
- Hernquist, L. 1990, *ApJ*, 356, 359
- Kaiser, N. 1987, *MNRAS*, 227, 1
- Khosroshahi, H. G., Ponman, T. J., & Jones, L. R. 2007, *MNRAS*, 377, 595
- Kneib, J.-P. et al. 2003, *ApJ*, 598, 804
- Koranyi, D. M. & Geller, M. J. 2000, *AJ*, 119, 44
- Ledlow, M. J., Voges, W., Owen, F. N., & Burns, J. O. 2003, *AJ*, 126, 2740
- Lemze, D., Broadhurst, T., Rephaeli, Y., Barkana, R., & Umetsu, K. 2009, *ApJ*, 701, 1336
- Lin, Y., Mohr, J. J., & Stanford, S. A. 2004, *ApJ*, 610, 745
- Lokas, E. L. & Mamon, G. A. 2003, *MNRAS*, 343, 401
- Lopes, P. A. A., de Carvalho, R. R., Kohl-Moreira, J. L., & Jones, C. 2009, *MNRAS*, 392, 135
- Mahdavi, A., Böhringer, H., Geller, M. J., & Ramella, M. 2000, *ApJ*, 534, 114
- Mahdavi, A. & Geller, M. J. 2001, *ApJ*, 554, L129
- Mahdavi, A., Geller, M. J., Böhringer, H., Kurtz, M. J., & Ramella, M. 1999, *ApJ*, 518, 69
- Mahdavi, A., Trentham, N., & Tully, R. B. 2005, *AJ*, 130, 1502
- Majumdar, S. & Mohr, J. J. 2004, *ApJ*, 613, 41
- Mandelbaum, R., Seljak, U., Cool, R. J., Blanton, M., Hirata, C. M., & Brinkmann, J. 2006, *MNRAS*, 372, 758

- Mantz, A., Allen, S. W., Ebeling, H., & Rapetti, D. 2008, *MNRAS*, 387, 1179
- Masson, C. R. 1979, *MNRAS*, 187, 253
- Merchán, M. E. & Zandivarez, A. 2005, *ApJ*, 630, 759
- Miller, C. J. et al. 2005, *AJ*, 130, 968
- Moore, B., Quinn, T., Governato, F., Stadel, J., & Lake, G. 1999, *MNRAS*, 310, 1147
- Mulchaey, J. S. & Zabludoff, A. I. 1998, *ApJ*, 496, 73
- Nagamine, K. & Loeb, A. 2003, *New Astronomy*, 8, 439
- Navarro, J. F., Frenk, C. S., & White, S. D. M. 1997, *ApJ*, 490, 493
- Osmond, J. P. F. & Ponman, T. J. 2004, *MNRAS*, 350, 1511
- Osmond, J. P. F., Ponman, T. J., & Finoguenov, A. 2004, *MNRAS*, 355, 11
- Ponman, T. J., Bourner, P. D. J., Ebeling, H., & Böhringer, H. 1996, *MNRAS*, 283, 690
- Popesso, P., Biviano, A., Böhringer, H., & Romaniello, M. 2007, *A&A*, 461, 397
- Popesso, P., Biviano, A., Böhringer, H., Romaniello, M., & Voges, W. 2005, *A&A*, 433, 431
- Popesso, P., Böhringer, H., Brinkmann, J., Voges, W., & York, D. G. 2004, *A&A*, 423, 449
- Quintana, H. & Melnick, J. 1982, *AJ*, 87, 972
- Rasmussen, J., Ponman, T. J., Mulchaey, J. S., Miles, T. A., & Raychaudhury, S. 2006, *MNRAS*, 373, 653
- Regös, E. & Geller, M. J. 1989, *AJ*, 98, 755
- Reisenegger, A., Quintana, H., Carrasco, E. R., & Maze, J. 2000, *AJ*, 120, 523
- Rines, K., Geller, M. J., Diaferio, A., Mahdavi, A., Mohr, J. J., & Wegner, G. 2002, *AJ*, 124, 1266
- Rines, K., Geller, M. J., Diaferio, A., Mohr, J. J., & Wegner, G. A. 2000, *AJ*, 120, 2338
- Rines, K., Geller, M. J., Kurtz, M. J., & Diaferio, A. 2003, *AJ*, 126, 2152
- Rines, K., Mahdavi, A., Geller, M. J., Diaferio, A., Mohr, J. J., & Wegner, G. 2001, *ApJ*, 555, 558
- Rines, K. J. & Diaferio, A. 2006, *AJ*, 132, 1275
- Rykoff, E. S. et al. 2008, *ApJ*, 675, 1106
- Serra, A. L., Angus, G. W., & Diaferio, A. 2009, *ArXiv e-prints*
- Slinglend, K., Batuski, D., Miller, C., Haase, S., Michaud, K., & Hill, J. M. 1998, *ApJS*, 115, 1
- Small, T. A., Sargent, W. L. W., & Hamilton, D. 1997, *ApJS*, 111, 1
- Smith, R. J., Lucey, J. R., Hudson, M. J., Schlegel, D. J., & Davies, R. L. 2000, *MNRAS*, 313, 469
- Stanek, R., Evvard, A. E., Böhringer, H., Schuecker, P., & Nord, B. 2006, *ApJ*, 648, 956
- Stoughton, C. et al. 2002, *AJ*, 123, 485
- Strauss, M. A. et al. 2002, *AJ*, 124, 1810
- Vedel, H. & Hartwick, F. D. A. 1998, *ApJ*, 501, 509
- Vikhlinin, A., McNamara, B. R., Forman, W., Jones, C., Quintana, H., & Hornstrup, A. 1998, *ApJ*, 502, 558
- Vikhlinin, A., McNamara, B. R., Hornstrup, A., Quintana, H., Forman, W., Jones, C., & Way, M. 1999, *ApJ*, 520, L1
- Voevodkin, A., Borozdin, K., Heitmann, K., Habib, S., Vikhlinin, A., Mescheryakov, A., & Hornstrup, A. 2009, *ArXiv e-prints*
- Voges, W. et al. 1999, *A&A*, 349, 389
- Wu, X.-P., Xue, Y.-J., & Fang, L.-Z. 1999, *ApJ*, 524, 22
- Xue, Y.-J. & Wu, X.-P. 2000, *ApJ*, 538, 65